



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SALVADOR.

No yellow fever in San Salvador.

SAN SALVADOR, *October 1, 1899.*

SIR: There does not exist a single case of yellow fever in San Salvador.

Respectfully, yours,

JOHN JENKINS,
United States Consul.

The SURGEON GENERAL,
U. S. Marine-Hospital Service.

FOREIGN STATISTICAL REPORTS.

AFRICA—*Lorenzo Marques*.—Month of August, 1899. Estimated population, 5,000. Total number of deaths, 25, including 2 from phthisis pulmonalis.

ARABIA—*Aden*.—Month of August, 1899. Estimated population, 41,346. Total number of deaths, 92. No deaths from contagious diseases reported.

BAHAMAS—*Dunmore Town*.—Two weeks ended October 13, 1899. Estimated population, 1,472. No deaths and no contagious diseases.

Governors Harbor.—Two weeks ended October 14, 1899. Estimated population, 1,500. Total number of deaths, 1. No contagious diseases reported.

Green Turtle Cay—Abaco.—Two weeks ended October 12, 1899. Estimated population, 3,900. No deaths and no contagious diseases.

Nassau.—Two weeks ended October 2, 1899. Estimated population, 12,000. Number of deaths not reported. No contagious diseases reported.

BRITISH GUIANA—*Demerara—Georgetown*.—Month of September, 1899. Estimated population, 36,567. Total number of deaths, 169. No contagious diseases reported.

CANADA—*Province of Ontario*.—Reports to the provincial board of health for the month of August from 675 localities having an aggregate estimated population of 2,225,326, show a total of 2,088 deaths, including diphtheria, 25; enteric fever, 55; measles, 5; scarlet fever, 8; whooping cough, 16, and 172 from phthisis pulmonalis.

FRANCE—*Roubaix*.—Month of September, 1899. Estimated population, 126,794. Total number of deaths, 188, including enteric fever, 1; measles, 4, and 3 from whooping cough.

GERMANY—*Glauchau*.—Month of September, 1899. Estimated population, 27,000. Total number of deaths, 62, including 4 from phthisis pulmonalis.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended October 7, 1899, correspond to an annual rate of 18.8 a thousand of the aggregate

population, which is estimated at 11,404,408. The highest rate was recorded in Salford, viz, 25.3, and the lowest in Swansea, viz, 12.1.

London.—One thousand five hundred and sixty-four deaths were registered during the week, including measles, 21; scarlet fever, 7; diphtheria, 55; whooping cough, 10; enteric fever, 19, and diarrhea and dysentery, 53. The deaths from all causes correspond to an annual rate of 17.9 a thousand. In Greater London 2,075 deaths were registered, corresponding to an annual rate of 16.6 a thousand of the population. In the "outer ring" the deaths included 10 from diphtheria, 5 from measles, 3 from scarlet fever, and 7 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended October 7, 1899, in the 22 principal town districts of Ireland was 24.7 a thousand of the population, which is estimated at 1,053,188. The lowest rate was recorded in Carrickfergus and Sligo, viz, 0.0, and the highest in Wexford, viz, 31.6 a thousand. In Dublin and suburbs 196 deaths were registered, including enteric fever, 4; measles, 32, and whooping cough, 2.

Scotland.—The deaths registered in 8 principal towns during the week ended October 7, 1899, correspond to an annual rate of 17.5 a thousand of the population, which is estimated at 1,587,414. The lowest mortality was recorded in Paisley, viz, 12.8, and the highest in Dundee, viz, 18.8 a thousand. The aggregate number of deaths registered from all causes was 534, including diphtheria, 6; measles, 11; scarlet fever, 8, and whooping cough, 5.

JAMAICA—Kingston.—Month of September, 1899. Total number of deaths, 91, including diphtheria, 1, and 16 from phthisis pulmonalis.

SPAIN—Valencia.—Two weeks ended October 10, 1899. Estimated population, 143,861. Number of deaths not reported. Eleven deaths from typhus.

SWITZERLAND.—Reports for the week ended September 9, 1899, from 15 cities and towns having an aggregate estimated population of 670,000, show a total of 228, including diphtheria, 4; enteric fever, 4; whooping cough, 5, and 25 from phthisis pulmonalis.